

## **Post-Operative Instructions**

### **Knee Arthroscopy, ACL Reconstruction with Bone-Patellar Tendon-Bone Autograft and Meniscal Repair**

#### **Day of surgery**

- A. Diet as tolerated
- B. Icing is important for the first 5-7 days post-op. While the post-op dressing is in place, icing should be done continuously. Once the dressing is removed on the third post-operative day, ice is applied for 20-minute periods 3-4 times per day. Care must be taken with icing to avoid frostbite. Alternatively, Cryocuff or Game-ready ice cuff can be used as per instructions.
  
- C. Pain medication as needed every 4-6 hours (refer to pain medication sheet).
- D. Make sure you have a physical therapy post-op appointment scheduled during the first week after surgery

#### **First Post-Operative Day**

- A. Continue icing
- B. Pain medication as needed

#### **Second Post-Operative Day Until Return Visit**

- A. Continue ice pack as needed.
- B. Unless otherwise noted, weight-bearing is toe-touch only for 4 weeks. At 4 weeks, you can bear as much weight on the affected leg as you can tolerate. Most patients use crutches for the first 2-3 weeks.
- C. Call our office @ 626-517-0022 to confirm your first postoperative visit, which is usually about 1-2 weeks after surgery. If you are experiencing any problems, please call our office or contact us via the internet at [pasadenaorthopedics.com](http://pasadenaorthopedics.com).

#### **Third Post-Operative Day**

- A. You may remove surgical bandage and shower this evening. Apply 4x4 (or similar size) waterproof bandage to these wounds prior to showering and when showering is complete apply fresh waterproof bandage. Please ensure that the bandage is large enough to completely cover the incision. You will need to follow this routine for 2 weeks after surgery.

## Rehabilitation Protocol: Combined Anterior Cruciate Ligament (ACL) Reconstruction and Meniscal Repair

Name: \_\_\_\_\_

Date: \_\_\_\_\_

Diagnosis: \_\_\_\_\_

Date of Surgery: \_\_\_\_\_

**Phase I (Weeks 0-4)**

- **Weightbearing:** Partial weight bearing (50%) as tolerated with brace locked in full extension
- **Hinged Knee Brace:**
  - Locked in full extension for ambulation and sleeping (**Weeks 0-1**)
  - Unlocked for ambulation and removed while sleeping (**Weeks 1-4**)
- **Range of Motion:** No flexion greater than 90 degrees
- **Therapeutic Exercises**
  - No deep bends for first 4 months
  - Quad/Hamstring sets and heel slides
  - Non-weightbearing stretch of the Gastroc/Soleus
  - Straight-Leg Raise with brace in full extension until quad strength prevents extension lag
  - **No Hamstring Stretching Until 4 Weeks Post-op**

**Phase II (Weeks 4-12)**

- **Weightbearing:** As tolerated – discontinue crutch use
- **Hinged Knee Brace:** Discontinue brace use when patient has achieved full extension with no evidence of extension lag
- **Range of Motion:** Maintain full knee extension – work on progressive knee flexion
- **Therapeutic Exercises**
  - No deep bends for first 4 months
  - Closed chain extension exercises
  - Hamstring stretching, toe raises, balance exercises
  - Progress to weightbearing stretch of the Gastroc/Soleus
  - Begin use of the stationary bicycle

**Phase III (Months 3-8)**

- **Weightbearing:** Full weightbearing
- **Range of Motion:** Full/Painless ROM
- **Therapeutic Exercises**
  - No deep bends for first 4 months
  - Begin hamstring strengthening
  - Advance closed chain strengthening exercises, proprioception activities
  - Begin use of the Stairmaster/Elliptical
  - **Can Start Straight Ahead Running at 3 Months**



**Aaron M. Gipsman, MD**  
Orthopedic Surgery  
Sports Medicine  
626-517-0022

**Phase IV (Months 8-10)**

- Continue with strengthening (quad/hamstring) and flexibility
- Begin cutting exercises and sport-specific drills
- Maintenance program for strength and endurance
- **Return to sports at 10-12 months**

**Comments:**

**Frequency:** \_\_\_\_ times per week

**Duration:** \_\_\_\_ weeks

**Signature:** \_\_\_\_\_

**Date:** \_\_\_\_\_